



Diabetes DIALOGUES

Less is More: The Promise of Weekly Insulin in Type 2 Diabetes

This activity is jointly provided by



acofp | AMERICAN COLLEGE
OF OSTEOPATHIC
FAMILY PHYSICIANS

This activity is supported by an educational grant from Lilly.

Co-Chairs



**Jay H Shubrook,
DO, FACOF, FACAFP**

Professor, Diabetologist
Department of Clinical Sciences and
Community Health
Touro University California, College of
Osteopathic Medicine
Vallejo, CA, USA



Carol H Wysham, MD

Clinical Professor of Medicine
University of Washington
Medical Manager, Diabetes and
Endocrinology Clinic
MultiCare Rockwood Clinic
Spokane, WA, USA

Activity Overview

Target Audience

This activity is intended for PCPs and other members of the healthcare team in the U.S. who care for patients with T2D.

Educational Objectives

After completing this activity, the participant should be better able to:

- Describe the place of novel insulin therapies, including once-weekly basal insulin, in treating type 2 diabetes
- Develop strategies for initiating, dosing, and titrating once-weekly basal insulin for type 2 diabetes
- Examine the consequences of delaying initiation and intensification of basal insulin in eligible patients
- Communicate to your patients the advantages of once-weekly basal insulin, considering patient concerns about weight gain and hypoglycemia

Agenda

- Once-Weekly Insulin: A Game Changer in a Crowded Market
- Practices for Initiating and Dosing Once-Weekly Basal Insulin in T2D
- The Promise of Once-Weekly Insulin

Accreditation Information



In support of improving patient care, Medical Learning Institute Inc is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Osteopathic Physician Continuing Medical Education

The American College of Osteopathic Family Physicians (ACOFP) is accredited by the American Osteopathic Association (AOA) to provide osteopathic continuing medical education for physicians.

ACOFP designates this live activity for a maximum of 1.0 AOA Category 1-A credits and will report continuing medical education (CME) credits commensurate with the physician's participation in this program.

Allopathic Physician Continuing Medical Education

Medical Learning Institute Inc (MLI) designates this live activity for a maximum of 1.0 *AMA PRA Category 1 Credit™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

MOC Statement



Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit. Participation information will be shared through the ACCME's Program and Activity Reporting System (PARS).

Disclosures



Disclosure & Conflict of Interest Policies

ACOFP is committed to providing learners with high-quality CME/CE activities that promote improvements in health care and not those of an ineligible company. In accordance with the ACCME Standards for Integrity and Independence in Accredited Continuing Education, also adopted by the AOA, ACOFP requires that individuals in a position to control the content of an educational activity disclose all relevant financial relationships with any ineligible company (those whose primary business is producing, marketing, selling, re-selling, or distributing healthcare products used by or on patients). Individuals in a position to control the content may include, but are not limited to, planning committee members, authors, faculty, speakers, reviewers, and activity staff.

Medical Learning Institute Inc is committed to providing high quality continuing education to healthcare professionals, as individuals and teams, with a protected space to learn, teach, and engage in scientific discourse free from influence from ineligible companies that may have an incentive to insert commercial bias into education. To that end, MLI requires faculty, presenters, planners, staff, and other individuals who are in a position to control the content of this CE activity to disclose all financial relationships they have had in the past 24 months with ineligible companies as defined by the ACCME, as related to the content of this CE activity, regardless of the amount or their view of the relevance to the education. All identified COI will be thoroughly vetted and mitigated according to MLI policy. These disclosures will be provided to learners prior to the start of the CE activity.

Disclosure of Unlabeled Use

This educational activity may contain discussions of published and/or investigational uses of agents that are not indicated by the FDA. The planners of this CE activity do not recommend the use of any agent outside of the labeled indications. The opinions expressed in the CE activity are those of the presenters and do not necessarily represent the views of the planners. Please refer to the official prescribing information for each product for discussion of approved indications, contraindications, and warnings.

Disclaimer

Participants have an implied responsibility to use the newly acquired information to enhance patient outcomes and their own professional development. The information presented in this CE activity is not meant to serve as a guideline for patient management. Any procedures, medications, or other courses of diagnosis or treatment discussed or suggested in this CE activity should not be used by clinicians without evaluation of their patient's conditions and possible contraindications and/or dangers in use, review of any applicable manufacturer's product information, and comparison with recommendations of other authorities.

Method of Participation

There are no fees for participating in or receiving credit for this CE activity. In order to receive credit, learners must participate in the entire CE activity, complete the activity evaluation form and your certificate of credit will be generated. A passing score of 70% or higher is needed to obtain credit. You will receive your certificate from Medical Learning Institute Inc.

For information on applicability and acceptance of continuing education credit for this CE activity, please consult your professional licensing board.

To submit questions to the presenter please send them to elearning@acofp.org and include the course title so we can direct them correctly.

If you have questions regarding your certificate, please contact MLI at mvu@mlieducation.org.

For Physician, If requesting MOC credit, the post-test and evaluation are required in their entirety as well as your ABIM ID number, DOB (MM/DD), and a score of 70% or higher is needed to obtain MOC credit.


Once-Weekly Insulin: A Game Changer in a Crowded Market

Treatment Challenges

-  Dose administration, timing, multiple injections
-  Hypoglycemia
-  Frequent dose adjustments
-  High number of injections (365/year)
-  Poor adherence
-  Weight gain/fluid retention
-  Only ~5% to 45% of patients achieve desired glycemic goals in the real world

69% to 87%

of people with type 2 diabetes (T2D) reported missing basal insulin dose injections



3.6 to 4.3 missed doses in a month significantly impacted glucose control



5.7 to 6.2 mistimed basal doses in a month significantly impact glycemic control

Opportunities of Weekly Basal Insulins



Advantages:


- Extend the time-action profile to allow for once-weekly administration
- Development of a once-weekly basal insulin with a longer, flatter exposure profile
- More closely mimic endogenous basal insulin distribution profiles
- Controlled tissue distribution properties and attenuated potency at the insulin receptor (IR)
- Reduce variability by controlling fluctuations in glucose levels during the week
- Maintaining an acceptable and manageable hypoglycemia profile



Limitations: inability to rapidly adapt to changes in insulin requirements, which the body achieves with controlled endogenous insulin secretion

However, patient preference for fewer injections may improve acceptance, adherence, and persistence

Fewer Injections = Improved Adherence



Once-weekly formulations of GLP-1 RAs are associated with significantly better adherence than once-daily formulations

Similar benefits may be observed with once-weekly basal insulins

Used with permission from Dr. Wysham.

^a Good adherence, $\geq 80\%$ proportion of days covered.

GLP-1 RA, glucagon-like peptide-1 receptor agonist; OD, once daily; OW, once weekly.

Qiao Q, et al. *Diabetes Metab Syndr Obes.* 2016;9:201-205.

Social Listening: Once-Weekly Insulin Concerns & Importance of Education

I would worry about any weekly medication. If you have a bad reaction to it, you are stuck with it for a week.

~DD



acofp

Social Listening: Once-Weekly Insulin Concerns & Importance of Education

A weekly insulin if they could make it, would be one heck of a HUGE shot too though! Imagine someone who needs 100 units a day, that'd be a 700 unit shot, egads! You'd look like you had a golf ball under your skin lol.

~RH



acofp

Habitual Prescribing?

- Proven efficacy in managing blood glucose
- Guideline-driven prescribing and standard of care
- Patient convenience
- Cost and insurance coverage: biosimilar options and insurance formularies
- Positive impact on comorbidities
- Familiarity to both physicians and patients

Addressing Hypoglycemia Concerns



How long would an episode of hypoglycemia last?

- Two-period crossover study in patients with T2D on insulin $\pm$ oral glucose-lowering meds
- Compared clinical, physiological, and counterregulatory hormone responses to 2 $\times$ & 3 $\times$ doses of icodec with insulin glargine U100
- 2 $\times$ & 3 $\times$ doses of once-weekly icodec do not lead to increased risk of hypoglycemia
- Time to develop hypoglycemia and recover were comparable



Would the episode recur?

- Continuous glucose monitoring (CGM) data: time spent in hypoglycemia in the weeks following the 2 $\times$ & 3 $\times$ doses was low
- Even in pts with clinically significant hypoglycemia (mean $\pm$ SD TBR [<54 mg/dL]: 2x dose $0.21 \pm .45\%$; 3x dose $0.56 \pm 1.70\%$)




How are acute episodes managed?

- Fundamentally, no differently than with once-daily basal insulins
- Administer calculated amounts of carbohydrates, monitor response, and repeat as necessary

CGM as Part of the Team

Compared to SMBG, rt-CGM is highly likely to be a cost-effective intervention for patients living with insulin-treated T2D



- The lower the mean patient age, the more cost-effective rt-CGM was likely to be
- CGM should be considered as an alternative to drug therapy for improving blood glucose

In 6-week intervention of of CGM with telemonitoring versus enhanced usual care:

	CGM	Enhanced Usual Care
Mean change HbA _{1C} , %	-0.69	-0.33
Body weight, kg	-2.48	+0.31

Examples of CGM Devices




FreeStyle Libre



Guardian Connect



Eversense



G6



G7

Social Listening: Patient Preference & Convenience

I was really happy that I only had to do an injectable once a week rather than at every meal or once a day. I was just amazed how something could stay in your system for a week and work for the whole week.

-Patient enrolled in ONWARDS-5
(insulin-naïve T2D)



acofp

Key Takeaways



Once-weekly basal insulins are promising options that address many of the barriers associated with once-daily formulations




Incorporating once-weekly basal insulins into T2D treatment plans will require shared decision-making supported by empowering, patient-first communication strategies, addressing individual patient needs/concerns, and detailed education

Practices for Initiating and Dosing Basal Insulin in T2D

Implications of Delayed Initiation

- Due to delays in initiation and intensification, only a minority of patients reach glycemic targets
- In the first 3 months, approximately 50% of patients interrupt their basal insulin therapy, and 15% discontinue it
- In patients with a delay in intensification of more than 1 year:
 - Reduced QOL
 - Increased risk of morbidity and mortality, including higher rates of myocardial infarction (67%), heart failure (64%), and stroke (51%)

Once-Weekly vs Daily Insulin



*Schematic representation of single doses

Initial Dosing



Learning Point!


Estimated starting basal insulin dose = 10 units/day OR 0.1 to 0.2 units/kg body weight/day
(depending on the degree of hyperglycemia)



Doses should be individualized and titrated over following days and weeks as needed

Fasting Plasma Glucose (FPG)	2 or more values of < 70 mg/dL	71 to 130 mg/dL	No values of < 70 mg/dL and 3 or more values > 130 mg/dL	No values of < 70 mg/dL and 3 or more values > 180 mg/dL
Insulin dose	Subtract 4 units	Maintain dose	Add 2 units	Add 4 units
Dose should be up/down titrated 1 to 2 times per week				

Clinical Signals That May Indicate Overbasalization



Blood Glucose Monitoring

People who take insulin should be encouraged to check their blood glucose levels



SMBG is essential for:

- Monitoring safety and efficacy of current treatment plan/dosing
- Hypoglycemia prevention and prompt treatment



Assessing FPG with SMBG to inform dose adjustments results in lower A1C levels



Increasingly, people with T2D have access to CGM and FGM, which encourages better awareness of blood glucose levels

Physician-Driven vs Self-Titration

The self-titration algorithm instructed participants to adjust their basal insulin dose every 3 days based on the mean of 3 adjusted FPG values

< 80 mg/dL

80 to 110 mg/dL


> 110 mg/dL

Reduce by
3 units

No change

Increase by
3 units

Physician-driven adjustments compared with self-titration algorithm-driven adjustments



■ Self-titration algorithm ■ Physician-driven adjustments

The Promise of Once-Weekly Insulin

Once-weekly basal insulin formulations are not currently approved by the FDA.

Advantages of Once-Weekly Insulin

Clinical	Molecular
Improved or similar glycemic control with low hypoglycemia risk	Long half-life
Reduced treatment burden	More stable pharmacokinetics/ pharmacodynamics, with less inter-patient and intra-patient variability
Easier to overcome therapeutic inertia	Slower clearance



Better treatment acceptance and adherence

In Development: Weekly Basal Insulin

Efsitora

QWINT Trial Program:


Topline Results of Phase III Trials

Icodec

ONWARDS Trial Program:

Results of 6 Phase III Trials

Insulin Efsitora: Efficacy and Safety of the Phase III QWINT Program





Learning Point!

Hypoglycemia event rates¹

1. Events per participant-year of exposure; 2. 95% CI, -0.22 to 0.04; 3. 95% CI, 0.94 to 1.78; 4. 95% CI, -0.22 to 0.061; 5. 95% CI, -0.19 to 0.12; 6. 95% CI, -0.22 to 0; 7. 95% CI, -0.075% to 0.19%.
 QWINT 1. <https://www.clinicaltrials.gov/ct2/show/NCT05662332>; QWINT 2. <https://www.clinicaltrials.gov/ct2/show/NCT05362058>; QWINT 3. <https://www.clinicaltrials.gov/ct2/show/NCT05275400>; QWINT 4. <https://www.clinicaltrials.gov/ct2/show/NCT05462756>;
 Eli Lilly and Company. With Once-a-Week Dosing, Insulin Efsitora Alfa Delivers A1C Reduction and Safety Profile Consistent with Daily Insulin. May 16, 2024; Eli Lilly and Company. In a first-of-its-kind fixed dose study, once weekly insulin efsitora alfa leads to A1C reduction similar to daily insulin. September 5, 2024; Bergenstal RM, et al. *Lancet*. 2024;404(10458):1132-1142; Wysham C, et al. *N Engl J Med*. 2024;391(23):2201-2211.

Insulin Icodec: Efficacy and Safety of the Phase III ONWARDS Program



In people with type 2 diabetes: No statistical difference in estimated hypoglycaemia events

*P<0.001 (noninferiority);
P=0.02 (superiority)

■ Once-weekly insulin icodec ■ Once-daily insulin glargine U100 ■ Once-daily insulin degludec ■ Once-daily basal insulins

*Statistically significant. 1 Severe or clinically significant hypoglycaemia events (blood glucose <3 mmol/L) per patient year, included for end of trial/end main phase in-trial. 2 Duration refers to trial main phase. ONWARDS 1: QW insulin icodec vs QD insulin glargine U100 both with non-insulin anti-diabetic treatment in insulin-naïve people with T2D; ONWARDS 2: QW insulin icodec vs QD insulin degludec in people with T2D switching from a QD insulin; ONWARDS 3: QW insulin icodec vs QD insulin degludec in insulin-naïve people with T2D; ONWARDS 4: QW insulin icodec vs QD insulin degludec both with mealtime insulin in people with T2D treated with basal and bolus insulin; ONWARDS 5: QW insulin icodec vs QD basal insulin with an app providing dosing recommendation in insulin-naïve people with T2D; ONWARDS 6: QW insulin icodec vs QD insulin degludec both with mealtime insulin in people with T1D. T1D: Type 1 diabetes; T2D: Type 2 diabetes. Note: Overview refer to primary end-points in main phases of trials

In Brief...



acofp



Insulin efsitora demonstrated non-inferior glycemic control compared with daily insulins



Insulin efsitora demonstrated a favorable safety profile in phase II clinical trials and topline results from the QWINT-2 and -4 phase III trials



Insulin icodec demonstrated superior glycemic control compared to daily insulins in 4/6 phase III ONWARDS clinical trials and non-inferiority in the other 2



Insulin icodec showed a favorable safety profile with no increased risk of hypoglycemia in patients with T2D



Once-weekly insulins offer a safe, effective option with a convenient dosing regimen, potentially reducing the burden of insulin therapy for persons with T2D



Reminder: Pework Activity & Clinical Data

Importance of Adherence in T2D

Impact on glucose control:

- 3.6 to 4.3 **missed** doses in a month significantly impacts glucose control
- 5.7 to 6.2 **mistimed** doses in a month significantly impacts glycemic control

Of people with T2D who reported missing basal insulin dose injections:

- 3× more likely to develop cardiovascular disease
- 10× more likely to develop kidney disease than people without diabetes
- 1/3 of people with diabetes will develop some form of vision loss during their lifetime
- In the USA, more than 100 000 people with diabetes have limbs amputated each year

Social Listening: Patient-First Language

Compliant. I really don't like that word. It is not like you are refusing the doctor's advice, you are just making the wrong decisions.

~BG



acofp

Social Listening: Non-Judgmental

Don't tell me I am a bad patient.
You don't have diabetes. You
don't know how hard it is. I try my
best to manage what I didn't ask
for.

~ EO



acofp

Preferences for Once-Weekly Basal Insulin

Patients are most concerned about:

- Avoiding 10-pound weight change
- Achieving longest time-in-range

Patient and Provider Preferences

Evaluating Patient and Provider Preferences for a Once-Weekly Basal Insulin in Adults with Type 2 Diabetes

A US survey-based study that included a discrete choice experiment among patients with T2D (insulin naïve and current insulin users) and HCPs who treat patients with T2D to assess preferences regarding a hypothetical once-weekly basal insulin in comparison to current basal insulin options.

Abbreviations: HCPs, healthcare providers; T2D, Type 2 diabetes; US, United States



Insulin-experienced patients with T2D were more confident that a once-weekly insulin would help with goals for blood glucose levels compared to their current basal insulin.



Social Listening: Dosing & Injection Challenges

Yesterday, I screwed up my basal injection and lost some of it, probably about half of it. I did not try to take any more because I didn't know how much I got.

~ KR



acofp

Social Listening: Weight Concerns


I've recently been put on insulin (Nov) and put on 28 lbs! It's making me more miserable than diabetes does – any advice on shifting insulin weight?

~KA



acofp

Communication and Collaboration



Learning Point!

- ✓ Upon diagnosis and whenever A1C is at or above goal: Discuss the **progressive nature** of T2D and the need to **frequently review and adjust treatments** in a timely manner
- ✓ **Discuss** people's concerns about **side effects and injections**
- ✓ Anticipate feelings of **guilt**
- ✓ Help people **anchor their treatment** to a daily activity
- ✓ Teach **self-titration** and provide **written instructions**; arrange **frequent contact** to ensure people are titrating
- ✓ Provide **feedback** based upon **glucose testing results**
 - Breaks a major link in the chain of clinical inertia
 - Promotes self-management

Talking Points and Top Tips



- Insulin is the most “natural” therapy, replacing what is missing (use thyroid hormone analogy)  Learning Point!
- Insulin has no drug interactions
- It prevents complications
- It can be successfully used in most people with minimal weight gain and hypoglycemia (especially when given early)



- Suggest people try insulin for 1 month and that they can stop if they do not feel it is helping
- If the clinic has a Diabetes Care & Education Specialist, can be used to demonstrate “dry shot” while learning glucose monitoring

In Brief...



Once-weekly basal insulin provides an opportunity for achieving patients' glycemic targets, reducing long-term complications of T2D, and improving patient adherence



Communication and collaboration can alleviate patient concerns about side effects and assist with patient education and long-term adherence